

APPROXIMATING CLASSIFYING SPACES BY SMOOTH PROJECTIVE VARIETIES

TORSTEN EKEDAH

ABSTRACT. We prove that for every reductive algebraic group H with centre of positive dimension and every integer K there is a smooth and projective variety X and an algebraic H -torsor $P \rightarrow X$ such that the classifying map $X \rightarrow BH$ induces an isomorphism in cohomology in degrees $\leq K$. This is then applied to show that if G is a connected non-special group there is a G -torsor $P \rightarrow X$ for which we do not have $[P] = [G][X]$ in the (completion of the) Grothendieck ring of varieties.

For a fixed integer k and a variable integer n , the Grassmannian $\mathrm{Gr}(k, n+k)$ of (complex) k -spaces in (complex) $n+k$ -dimensional space becomes a closer and closer (the larger we make n) homotopy approximation to the classifying space of $\mathrm{GL}_k(\mathbf{C})$ in the following sense: The frame bundle of the tautological k -bundle is a $\mathrm{GL}_k(\mathbf{C})$ -torsor over $\mathrm{Gr}(k, n+k)$ and thus gives a map $\mathrm{Gr}(k, n+k) \rightarrow \mathrm{BGL}_k(\mathbf{C})$ which induces an isomorphism on homotopy groups up to a degree which tends to infinity with n . The point here is that $\mathrm{Gr}(k, n+k)$ is a complex projective manifold and that the $\mathrm{GL}_k(\mathbf{C})$ -torsor is also algebraic. Hence, we can homotopically approximate the classifying space of $\mathrm{GL}_k(\mathbf{C})$ by projective manifolds using algebraic torsors. This is somewhat special to $\mathrm{GL}_k(\mathbf{C})$; it is for instance not possible to do it for $\mathrm{SL}_k(\mathbf{C})$ as $H^2(\mathrm{BSL}_k(\mathbf{C}), \mathbf{Z}) = 0$ and the second cohomology group of a projective manifold is never zero. When G is a finite group this was rectified by Atiyah and Hirzebruch (cf., [AH62, Prop. 6.6]) who showed that the Godeaux-Serre method (cf., [Se58]) together with the Lefschetz hyperplane theorem could be used to prove that we can get homotopy approximations to $\mathrm{B}(G \times \mathbf{G}_m)$ which are given by algebraic torsors over a smooth and projective base. The main result of the present paper is an extension of this result to the case when G is a reductive group (and where we also allow ourselves to replace $G \times \mathbf{G}_m$ by any central extension of G by $\mathbf{G}_m$).

The technical problems we have to fight with is the fact that when $\dim G > 0$ there will be unstable points for the action of G on a projective space. The unstable points will contribute to the cohomology of a (suitably iterated) hypersurface section defined by a G -invariant form which will have to be “subtracted off” if we are to get hold of the cohomology of the torsor (and consequently its smooth projective base). This requires us to get information on the cohomology of the unstable locus and we shall use the inductive analysis of the unstable points introduced by Kempf et al. We are much helped by the fact that we can allow ourselves to consider $\mathbf{P}(V^n)$ where V is a fixed linear representation and n is arbitrarily large; making n large enough means that big pieces of the cohomology of the unstable locus will not interfere with our goal. On the other hand, in the argument we shall use (repeatedly) Poincaré duality and consequently our results will only give a homological and not a homotopy approximation.

The fact that our constructions are very non-canonical makes it unlikely that they could be used to study the the classifying spaces of reductive (or equivalently compact) groups. Instead it can be used to show the existence of projective manifolds exhibiting various topological behaviours (as was already demonstrated by Atiyah and Hirzebruch where a particular behaviour of cohomology under the action of the Steenrod algebra was required). Indeed, we shall apply it to show that if G is a linear connected non-special (in the sense of Grothendieck and Serre) group, then there are algebraic G -torsors with a smooth projective base such that the associated G/B -fibration has cohomology which is not (additively) isomorphic to the cohomology of the product of the base and G/B . (On the contrary such an isomorphism always exists for a special group.) This will then be used to obtain the result on the Grothendieck ring of varieties mentioned in the abstract.

1 The approximation

So far we have formulated our result in topological terms and that forces us to work with algebraic groups over the complex numbers. In Subsection 1.1 we shall discuss the necessary modifications to obtain results valid over any field.

For the proof of our first proposition we need to recall (cf., [Ki84, Part II] whose notation we shall follow) the inductive analysis of unstable points of a representation W of a reductive group G . Hence, if we fix a maximal torus T of G we have a stratification $\{S_\beta\}$ of $\mathbf{P}(W)$ parametrised by certain elements of $M \otimes \mathbf{Q}$, where M is the cocharacter group of T . To such a $\beta \neq 0$ the following data can be associated:

- A positive definite quadratic form q on $M \otimes \mathbf{Q}$ that is invariant under the Weyl group has been chosen once and for all.
- Putting $N_\beta := \{\alpha \in N \mid \alpha \cdot \beta = q(\beta)\}$ and $N'_\beta := \{\alpha \in N \mid \alpha \cdot \beta > q(\beta)\}$, where N is the character group of T , we let $W_\beta \subseteq W$ be the subspace spanned by the T -eigenvectors of weights belonging to N_β .
- We let T_β be the image of $r\beta$, where r is a strictly positive integer such that $r\beta \in M$. The centraliser Stab_β of T_β is a reductive group stabilising W_β . There is a reductive subgroup G_β of Stab_β such that $\text{Stab}_\beta = T_\beta G_\beta$ and T_β is not contained in G_β . In particular $\dim \text{Stab}_\beta = \dim G_\beta + 1$.
- Setting $Z_\beta := \mathbf{P}(W_\beta) \subseteq \mathbf{P}(W)$ we have that Stab_β acts (linearly) on it. We let $Z_\beta^{ss} \subseteq Z_\beta$ be the semi-stable locus for the action of G_β .
- Letting $W'_\beta \subseteq W$ be the subspace spanned by the T -eigenvectors of weights belonging to N'_β , $W'_\beta + W_\beta \subseteq W$ is a direct sum. Putting $Y_\beta := \mathbf{P}(W_\beta \oplus W'_\beta) \setminus \mathbf{P}(W'_\beta)$ we have a linear projection morphism $Y_\beta \rightarrow Z_\beta$ which is a vector bundle of constant rank $r_\beta(V) := \dim W'_\beta$. (Under the identification of $Y_\beta \rightarrow Z_\beta$ with a vector bundle Z_β becomes the zero section.) We let Y_β^{ss} be the inverse image of Z_β^{ss} under the map $Y_\beta \rightarrow Z_\beta$.
- There is a parabolic subgroup P_β stabilising Y_β and Y_β^{ss} as well as containing Stab_β . More precisely Stab_β is a Levi factor of P_β , i.e., it maps bijectively to P_β/U_β , where U_β is the unipotent radical of P_β . In particular, as U_β and G/P_β have the same dimension we have that $\dim G = \dim \text{Stab}_\beta + 2 \dim(G/P_\beta)$.
- We have an isomorphism $S_\beta \cong G \times_{P_\beta} Y_\beta^{ss}$ and S_0 equals $\mathbf{P}(W)^{ss}$, the G -semistable locus of $\mathbf{P}(W)$. In particular if there is no weight of W that lies in N_β we have that $S_\beta = \emptyset$.

We now denote by S'_β , $Y_\beta'^{ss}$ and $Z_\beta'^{ss}$ the inverse images under the quotient map $W \setminus \{0\} \rightarrow \mathbf{P}(W)$ of S_β , Y_β^{ss} and Z_β^{ss} respectively. The linear projection map $Y_\beta \rightarrow Z_\beta$ induces a vector bundle map $Y_\beta'^{ss} \rightarrow Z_\beta'^{ss}$ of rank $r_\beta(W)$. Consequently we have that

$$H_c^i(Y_\beta'^{ss}, \mathbf{Z}) = H_c^{i-2r_\beta(W)}(Z_\beta'^{ss}, \mathbf{Z}) \quad (1)$$

and in particular $H_c^i(Y_\beta'^{ss}, \mathbf{Z}) = 0$ whenever $i < 2r_\beta(W)$. We also have that $S'_\beta = G \times_{P_\beta} Y_\beta'$. Its pullback along $G \rightarrow G/P_\beta$ is a product and as P_β is connected, the proper base change theorem implies that $R^i \pi_! \mathbf{Z}$ is constant with constant value $H_c^i(Y_\beta'^{ss}, \mathbf{Z})$. This (together with the fact that G/P_β is compact) implies that the Leray spectral sequence has the form

$$H^j(G/P_\beta, H_c^i(Y_\beta'^{ss}, \mathbf{Z})) \implies H_c^{i+j}(S'_\beta, \mathbf{Z}). \quad (2)$$

Turning now to a slightly different topic let V be a k -dimensional complex vector space for $k > 0$. We define the open subset $U_n \subseteq V^n$ by the condition that $(v_1, \dots, v_n) \in U_n$ if there is a subset $S \subseteq \{1, \dots, n\}$ with $|S| = k$ and such that $(v_s)_{s \in S}$ is a basis for V . Clearly, U_n is stable under multiplication by scalars and so corresponds to an open subset W_n of $\mathbf{P}(V^n)$.

Lemma 1.1 i) A tuple $(v_1, \dots, v_n)$ lies in U_n precisely when the v_i span V .
 ii) $\mathrm{GL}(V)$ acts freely on U_n and hence so does any algebraic subgroup of $\mathrm{GL}(V)$.
 iii) Any point of W_n is semi-stable (and therefore also stable by ii) for the action by $\mathrm{SL}(V)$ and thus also for any algebraic subgroup of $\mathrm{SL}(V)$.
 iv) The codimension of $V^n \setminus U_n$ in V^n is $\geq n + 1 - k$.
 v) Let $G \subseteq \mathrm{SL}(V)$ be a (not necessarily connected) reductive algebraic subgroup. Let $V_{ss}^n \subseteq V^n \setminus \{0\}$ be the inverse image of the locus of G -semistable points of $\mathbf{P}(V^n)$. Then for every K there is an integer N such that $H_c^i(V_{ss}^n, \mathbf{Z}) = 0$ for all $\dim G + 1 < i \leq K$ and $n \geq N$.

Proof. To begin with i) is just linear algebra.

For ii) a point $(v_1, \dots, v_n)$ of U_n contains a basis and hence its stabiliser is reduced to the identity. Hence it remains to show that the morphism $\mathrm{GL}(V) \times U_n \rightarrow U_n \times U_n$ given by $(g, x) \mapsto (x, gx)$ is proper. For this we use the valuation criterion so we assume that we have a discrete valuation ring R with fraction field K and K -points g and x such that x and gx are defined as R -points of U_n . As x and gx thus both span $V \otimes R$ we get that g and its inverse are defined over R .

As for iii), for every $S \subseteq \{1, \dots, n\}$ we have the $k \times S$ -determinant which is $\mathrm{SL}(V)$ -invariant and for every point of U_n there is an S such that it separates the point from the origin.

A point $(v_1, \dots, v_n)$ in the complement of U_n has, by i), the property that its linear span is a proper subspace of V . Hence, the point lies in the union of the U^n , where U runs over the hyperplanes of V . That union has dimension at most $k - 1 + n(k - 1)$ and hence codimension at least $nk - (n + 1)(k - 1) = n + 1 - k$ which proves iv).

Finally, to prove v) we start by noticing that if G° is the group of connected components of G , then a point of $\mathbf{P}(V^n)$ is semi-stable for G precisely when it is so for G° . Hence we may, to simplify, assume that G is connected.

We now apply the analysis recalled above of unstable points of the G -representation V^n . The crucial part is that by construction $(V^n)_\beta = (V_\beta)^n$ (as N_β does not depend on the representation). Furthermore, again by construction, the set of β for which the S_β are non-empty is contained in a finite set which only depends on the weights of T appearing in the V^n and is thus independent of n . We now prove the result by induction on $\dim G$. The base case is when $G = \{e\}$ in which case $V_{ss}^n = V^n \setminus \{0\}$ and thus $H_c^i(V_{ss}^n, \mathbf{Z}) = 0$ unless $i = 1$ or $i = 2n \dim V$ which gives the result in that case. In the general case, when $\dim G > 0$, we have that $\dim G_\beta < \dim G$ for all $\beta \neq 0$ so we may assume that the theorem is true for (G_β, V_β) . The complement S' of V_{ss}^n in $V^n \setminus \{0\}$ is the union of the S_β where $\beta \neq 0$ runs over a finite set, independent of n . Using the long exact sequences of cohomology with compact supports we can analyse the vanishing of $H_c^i(S', \mathbf{Z})$ in terms of the vanishing of the $H^i(S'_\beta)$. We have that $r_\beta(V^n) = nr_\beta(V)$ so if $r_\beta(V) \neq 0$ we get from (1) (and (2)) that by making n large enough we may assume that $H_c^i(S'_\beta, \mathbf{Z}) = 0$ when $i \leq K$. For those β with $r_\beta(V) = 0$ we have that $Y_\beta^{ss} = Z_\beta^{ss}$. Hence (2) and the induction assumption implies that by making n large enough we may assume that $H_c^i(S'_\beta, \mathbf{Z}) = 0$ whenever $2 \dim(G/P_\beta) + \dim G_\alpha + 1 < i \leq K$ and as, which was noted above, $2 \dim(G/P_\beta) + \dim G_\alpha + 1 = \dim G$ we get that $H_c^i(S'_\beta, \mathbf{Z}) = 0$ when $\dim G < i \leq K$. From this (and the long of exact sequences of cohomology) we get that $H_c^i(S', \mathbf{Z}) = 0$ for $\dim G < i \leq K$. We now have a long exact sequence

$$\dots \rightarrow H_c^{i-1}(V^n \setminus \{0\}, \mathbf{Z}) \rightarrow H_c^{i-1}(S', \mathbf{Z}) \rightarrow H_c^i(V_{ss}^n, \mathbf{Z}) \rightarrow H_c^i(V^n \setminus \{0\}, \mathbf{Z}) \rightarrow \dots$$

From it, and the fact that $H_c^i(V^n \setminus \{0\}, \mathbf{Z}) = 0$ for $1 < i < n \dim V$ it follows that $H_c^i(V_{ss}^n, \mathbf{Z}) = 0$ if $\dim G + 1 < i \leq K$ (and n is large enough). $\square$

The following result is no doubt well-known but lacking a reference we provide a proof.

Lemma 1.2 Let G be reductive group provided with a centrally embedded $\mathbf{G}_m \hookrightarrow G$. Then G has a faithful linear representation V such that the central $\mathbf{G}_m$ acts by the identity map into the scalar linear maps.

Proof. We first prove the following: Let $\mathcal{L}$ be a line bundle on an affine scheme X (over some affine base) and let R be the ring generated (over the base) by the sections of $\mathcal{L}$ considered as functions on the associated $\mathbf{G}_m$ -torsor T . Then the induced map $T \rightarrow \mathbf{Spec} R$ is an open embedding. Indeed, assume first that $\mathcal{L} = Rt$. In that case T maps into the subscheme $\mathbf{Spec} R[t^{-1}]$ but $R[t^{-1}]$ is equal to the affine coordinate ring of T so that $T \rightarrow \mathbf{Spec} R[t^{-1}]$ is an isomorphism. As $\mathcal{L}$ is locally trivial this implies that there is an open covering $\{U_i\}$ of T such that $U_i \hookrightarrow X \rightarrow \mathbf{Spec} R$ is an open embedding. From this it follows that it remains to prove that $X \rightarrow \mathbf{Spec} R$ is injective on points. That the elements of $\mathcal{L}$ separates points of $\mathcal{L}$ is clear however.

In the case of $\mathbf{G}_m \hookrightarrow G$ we get an $\mathcal{L}$ by considering the functions on G that are of weight 1 with respect to the $\mathbf{G}_m$ -action. It is a line bundle over $G/\mathbf{G}_m$. As $\mathbf{G}_m$ is central, $\mathcal{L}$ is G -invariant and is hence the union of finite dimensional subrepresentations. Applying what was just shown we get that a large enough such subrepresentation is faithful. This concludes our proof as by construction $\mathbf{G}_m$ acts on it by the identity map into scalars. $\square$

Before passing to our main theorem we want to emphasise that we do not assume that a reductive group is connected.

Theorem 1.3 *Let H be a reductive algebraic group with positive-dimensional centre over the complex numbers. Then for every integer K there is a smooth and projective variety X and an algebraic H -torsor $P \rightarrow X$ such that its classifying map $X \rightarrow BH$ induces an isomorphism $H^i(BH, \mathbf{Z}) \rightarrow H^i(X, \mathbf{Z})$ for all $i \leq K$.*

If H has the form $G \times \mathbf{G}_m$, then it is possible to choose P such that the line bundle associated to the $\mathbf{G}_m$ -torsor $P/G \rightarrow X$ is ample.

Proof. As the centre of H is positive dimensional and as it is reductive, the connected component of the centre is a torus so that we may find a copy of $\mathbf{G}_m$ contained in the centre. By Lemma 1.2 we may find a faithful linear representation V of G which restricts to multiplication by scalars on the $\mathbf{G}_m$. We now put $G := H \cap \mathrm{SL}(V)$. After possibly replacing V by V^n we may, by Lemma 1.1 and the condition that H acts faithfully on V , assume that there is an open subset $U \subseteq V$, whose complement has arbitrarily high codimension, stable under H such that H stabilises U and acts freely on it. Furthermore, again by the lemma, we may also assume that $U' := U/\mathbf{G}_m \subseteq \mathbf{P}(V)$ consists of G -stable points so that U is a H -torsor over an open subset U'' of the GIT quotient $\mathbf{P}(V)//G$. Finally, we may choose a suitable k such that the base point locus of $(S^k V)^G$ in $\mathbf{P}(V)$ is equal to the set T of unstable points and that $(S^k V)^G$ induces an embedding of the $\mathbf{P}(V)//G$ of $\mathbf{P}(V)$ by G into $\mathbf{P}((S^k V)^G)$.

We now construct a sequence $\mathbf{P}(V) = X_0 \supseteq X_1 \supseteq \dots \supseteq X_a$ of closed subvarieties such that

- the codimension of X_j in $\mathbf{P}(V)$ at any of its points outside of T is equal to i ,
- X_{j+1} is the zero-set in X_j of some $f_j \in (S^k V)^G$,
- X_j is smooth outside T and
- $X_a \setminus T$ lies inside U' .

That such a sequence exists is clear as the general hyperplane section of a linear system is smooth outside the base points of the system. Furthermore, as the codimension of $V \setminus U$ in V can be made arbitrarily large, once again by the lemma, so can L , the dimension of $X_a \setminus T$. We may also by (1.1:v) assume that $H_c^i(V \setminus T', \mathbf{Z}) = 0$ when $\dim H = \dim G + 1 < i < L$.

Now, by the fact that $X_j \setminus X_{j+1}$ is smooth and affine we get that $H_c^i(X_j, \mathbf{Z}) \rightarrow H_c^i(X_{j+1}, \mathbf{Z})$ is an isomorphism for $i < \dim(X_{j+1} \setminus T)$. Using the long exact sequence of cohomology (and the 5-lemma) this implies that $H_c^i(X_j \setminus T, \mathbf{Z}) \rightarrow H_c^i(X_{j+1} \setminus T, \mathbf{Z})$ is an isomorphism for $i < \dim(X_{j+1} \setminus T)$. Hence we get that $H_c^i(\mathbf{P}(V) \setminus T, \mathbf{Z}) \rightarrow H_c^i(X_a \setminus T, \mathbf{Z})$ is an isomorphism for $i < L$ which in turn implies that we have an isomorphism $H_c^i(V \setminus T', \mathbf{Z}) \rightarrow H_c^i(U'_a, \mathbf{Z})$, where U'_a is the inverse image of $X_a \setminus T$ in $V \setminus \{0\}$ and T' the inverse image of T . Thus we get that $H_c^i(U'_a) = 0$ for $\dim H < i < L$. Now, by construction H acts freely on U'_a and the quotient Y_a

is the zero set in $\mathbf{P}(V)//G$ of the f_i . Hence Y_a is proper as it is closed in $\mathbf{P}(V)//G$ and smooth as U'_a is. Furthermore, the Y_j are general iterated hyperplane sections and hence normal (by [Se50] and the fact that $\mathbf{P}(V)//G$ is) and therefore they are connected by the Lefschetz theorem (and of course the fact that $\mathbf{P}(V)//G$ is connected). This may be applied to the case when H is replaced by its connected component H^o and thus also U'_a is connected (as $U'_a \rightarrow U'_a/H^o$ is a fibration with connected fibres). Furthermore, if $K \subseteq H$ is a maximal compact subgroup, then the G -torsor $U'_a \rightarrow Y_a$ has a reduction to a K -torsor $Q_a \subseteq U'_a$ and this inclusion is a homotopy equivalence. Furthermore, as Y_a is a compact (complex) manifold, Q_a is also a compact manifold. Now, letting $K^o := H^o \cap K$ the fibration $Q_a \rightarrow Q_a/K^o$ has connected structure group and is hence orientable and as $Q_a/K^o = U'_a/G^o$ is a complex manifold and in particular orientable we conclude that Q_a is a compact orientable manifold.

We now apply duality to U'_a and conclude that $H_i(U'_a, \mathbf{Z}) = 0$ for $2d - L < i < 2d - h$, where d is the (complex) dimension of U'_a and $h := \dim H$. As Q_a is homotopic to U'_a the same is true for Q_a . Being an orientable manifold we can apply duality to it together with the fact that $\dim U'_a - \dim Q_a = \dim H - \dim K = h$. This gives $H^i(Q_a, \mathbf{Z}) = 0$ for $0 < i < L - h$ and as U'_a is connected so is Q_a and we get that $H^0(Q_a, \mathbf{Z}) = \mathbf{Z}$. Therefore we get that $H^i(U'_a, \mathbf{Z})$ equals $\mathbf{Z}$ if $i = 0$ and 0 if $0 < i < L - h$. We have a commutative diagram of H -torsors

$$\begin{array}{ccc} U'_a & \longrightarrow & EH \\ \downarrow & & \downarrow \\ Y_a & \longrightarrow & BH \end{array}$$

and the map $U'_a \rightarrow EH$ induces an isomorphism in cohomological degrees $< L - h$. This implies that the same is true for $Y_a \rightarrow BH$ (by for instance the spectral sequence $H^i(H^j \times X) \implies H^{i+j}(Y)$ that exists for any H -torsor $X \rightarrow Y$) and as L can be made arbitrarily large we are finished with the first part.

For the last part, we have by construction that $P/\mathbf{G}_m = X_a$ and thus $P \rightarrow P/\mathbf{G}_m$ is the $\mathbf{G}_m$ -torsor associated to the restriction $\mathcal{L}$ of $\mathcal{O}(1)$ to X_a and as G acts compatibly on P and $P/\mathbf{G}_m$, $\mathcal{L}$ extends to a line bundle $\mathcal{M}$ on $P/H = Y_a$. Furthermore, the space of sections of $\mathcal{M}^{\otimes r}$ equals $(S^r V)^G$ which shows that $\mathcal{M}^{\otimes k}$ is the restriction of the $\mathcal{O}(1)$ of $\mathbf{P}((S^k V)^G)$ to $Y_a \subseteq \mathbf{P}(V)//G \subseteq \mathbf{P}((S^k V)^G)$. $\square$

Remark: i) If H is connected its centre is positive-dimensional precisely when H is not semi-simple.

ii) For a general reductive group G , $G \times \mathbf{G}_m$ fulfils the conditions of the theorem.

iii) The proof of the theorem shows that we can refine (1.1:v) to $H_c^i(V_{ss}^n, \mathbf{Z}) = 0$ for all $0 < i \leq K$ and $i \neq \dim G + 1$ and $H_c^{\dim G + 1}(V_{ss}^n, \mathbf{Z}) = \mathbf{Z}$. It would be interesting to have a direct proof of this fact.

iv) Note that the appropriate homotopy version of the Lefschetz hyperplane theorem (cf., [Mi63, Thm 7.4]) gives that X_a is a homotopy approximation to $\mathbf{P}(V)$. However, we need to have information on the relation between $\mathbf{P}(V) \setminus T$ and $X_a \setminus T$. Our proof of such a relation involved the use of duality. Duality is an intrinsically stable (in the sense of homotopy theory) notion and hence destroys information on for instance the fundamental group. Note incidentally that T may very well have a larger dimension than $X_a \setminus T$.

However, it seems reasonable to believe that the restriction to homological approximation is just an artifice of the proof and I conjecture that there are G -torsors over smooth and projective varieties such that the classifying map induces an isomorphism on homotopy groups up to any limit.

v) One can, if one is so inclined, make an *a posteriori* use of the Lefschetz theorem to obtain a stronger result: Given an integer L one can find an algebraic H -torsor $P \rightarrow Y$ with Y smooth projective of dimension L such that $H^i(BH, \mathbf{Z}) \rightarrow H^i(Y, \mathbf{Z})$ is an isomorphism for $i < L$ and injective with a torsion free cokernel for $i = L$. Then $H^i(Y, \mathbf{Z})$ for $i > L$ is determined by duality.

One constructs Y from the X of the theorem (using a $K = L + 1$) by taking the appropriate number of hyperplane sections.

1.1 Algebraic version

In this subsection we shall quickly indicate the modifications needed to obtain a theorem valid over an arbitrary (algebraically closed, for simplicity) field $\mathbf{k}$, replacing of course classical cohomology with étale cohomology with values in $\widehat{\mathbf{Z}}'$, the inverse limit $\varprojlim_n \mathbf{Z}/n\mathbf{Z}$, where n runs over integers invertible in $\mathbf{k}$.

The classifying map $X \rightarrow BH$ of an H -torsor $\pi: P \rightarrow X$ makes sense if we replace the classifying space with the classifying algebraic stack. Apart from that, the only non-algebraic component of the proof is the use of the maximal compact subgroup K and the reduction of the structure group of a H -torsor to K . However, we only need the cohomological consequences of that fact. More precisely we need to know that the dual (when the base is smooth) of $R\pi_*\widehat{\mathbf{Z}}'$ is isomorphic to $R\pi_*\widehat{\mathbf{Z}}'[-h]$, where $h := \dim H$. For this we first note that the proof of the theorem shows that $R^i\pi_*\widehat{\mathbf{Z}}'$ is a constant sheaf of value $H^i(H, \widehat{\mathbf{Z}}')$. (Formally, the argument shows this only for direct images with compact support but we can use duality or smooth base change instead.) Suppose now that we know that $H^i(H, \widehat{\mathbf{Z}}') = 0$ for $i > h$, $H^h(H, \widehat{\mathbf{Z}}') = \widehat{\mathbf{Z}}'$ and that the multiplication map $R\Gamma(H, \widehat{\mathbf{Z}}') \otimes^L R\Gamma(H, \widehat{\mathbf{Z}}') \rightarrow R\Gamma(H, \widehat{\mathbf{Z}}')$ composed with $R\Gamma(H, \widehat{\mathbf{Z}}') \rightarrow H^h(H, \widehat{\mathbf{Z}}')[-h]$ gives an isomorphism $R\Gamma(H, \widehat{\mathbf{Z}}') \xrightarrow{\sim} R\mathrm{Hom}_{\widehat{\mathbf{Z}}'}(R\Gamma(H, \widehat{\mathbf{Z}}'), H^h(H, \widehat{\mathbf{Z}}'))[-h]$. We then get a morphism $R\pi_*\widehat{\mathbf{Z}}' \rightarrow R^h\pi_*\widehat{\mathbf{Z}}'[-h] = H^h(H, \widehat{\mathbf{Z}}')[-h]$ and it together with the multiplication map $R\pi_*\widehat{\mathbf{Z}}' \otimes^L R\pi_*\widehat{\mathbf{Z}}' \rightarrow R\pi_*\widehat{\mathbf{Z}}'$ gives a morphism $R\pi_*\widehat{\mathbf{Z}}' \rightarrow R\mathrm{Hom}_{\widehat{\mathbf{Z}}'}(R\pi_*\widehat{\mathbf{Z}}', H^h(H, \widehat{\mathbf{Z}}'))[-h]$. To check that this is isomorphism we can pull back by π and by smooth base change the map becomes just the constant map $R\Gamma(H, \widehat{\mathbf{Z}}') \xrightarrow{\sim} R\mathrm{Hom}_{\widehat{\mathbf{Z}}'}(R\Gamma(H, \widehat{\mathbf{Z}}'), H^h(H, \widehat{\mathbf{Z}}'))[-h]$.

Now, in order to show duality for $R\Gamma(H, \widehat{\mathbf{Z}}')$ we start by noticing that the cohomology of H is isomorphic to that of H/U , where U is the unipotent radical of a Borel group B of H . The quotient map $G/U \rightarrow G/B$ is a T -torsor and by the argument just given and the fact G/B is proper we are reduced to proving duality for $R\Gamma(T, \widehat{\mathbf{Z}}')$. This of course is well-known. (For the proof of duality for $R\Gamma(H, \widehat{\mathbf{Z}}')$ we could also have started off from the truth of it over $\mathbf{C}$ and then used a specialisation argument as in [De77, Sommes trig.:8.2] to show that the cohomology is independent of the algebraically chosen base field and in particular of the characteristic.)

In characteristic zero this gives us a purely algebraic proof of the theorem. In positive characteristic the problem is that we do not know that a general member of the linear system of H -invariant forms on $\mathbf{P}(V)$ is smooth outside of the base (i.e., unstable) locus. This can be rectified by going directly to (in the notations of the theorem) X_a : Letting $V_i := \{x \in \mathbf{P}(V) \mid f_i(x) \neq 0\}$ we have that the V_i as well as their intersections are smooth and affine and their union, V' , is the complement of X_a . Using the Čech spectral sequence for $V' = \cup_i V_i$ we get that $H^i(V', \widehat{\mathbf{Z}}') = 0$ for $i > \dim V' + a$ and using duality (as V' is smooth) and the long exact sequence of cohomology with compact support we get the needed properties of the cohomology of X_a .

Finally, as we no longer (in positive characteristic) can assume that $X_a \setminus T$ is smooth we can not use it to get the smoothness for X_a/H . However, X_a/H is a complete intersection with respect to a very ample linear system of $\mathbf{P}(V)//G$ lying in the smooth locus of $\mathbf{P}(V)//G$ so that if the f_i are chosen to be general the smoothness of X_a/H follows directly.

We can go even further and start with an algebraic group over an arbitrary (i.e., non-algebraically closed) field. We can then choose our representation V to be defined over the same field. The genericity conditions for the choice of the H -invariant forms can be fulfilled over the base field if it is infinite. In the case of a finite base field it can be fulfilled by increasing the degree of the forms.

2 Applications

An example of how the theorem can be applied to finite groups is given in [Ek86]. There one uses the fact that for every prime p there is a finite group G with elements $x, y, z \in H^1(G, \mathbf{Z}/p\mathbf{Z})$ such that the Massey product $\langle x, y, z \rangle$ is defined and non-trivial. Using a smooth and projective approximation one concludes that there is a smooth and projective variety with non-trivial $\mathbf{Z}/p\mathbf{Z}$ -Massey products (on the contrary rational Massey products of smooth and proper varieties are always trivial, cf., [DGMS]).

A somewhat vaguer application is to the theory of characteristic classes of G -torsors. The theorem implies that a characteristic class of G -torsors is determined by its restriction to the category of smooth projective varieties and algebraic G -torsors for G a reductive group.

Recall (cf., [SC58]) that an algebraic group G (over an algebraically closed field) is said to be *special* if every algebraic G -torsor is trivial in the Zariski topology. Special groups have the property that for every (topological) G -torsor $P \rightarrow X$, the associated G/B -fibration $G/B \times_G P =: Y \rightarrow X$ has $H^*(Y)$ isomorphic to $H^*(G/B \times X, \mathbf{Z})$. We can use the theorem to prove a very specific version of the converse of this result.

Proposition 2.1 *Let G be a connected non-special linear group. There is a smooth projective variety X and an algebraic G -torsor $P \rightarrow X$ such that if $Y \rightarrow X$ is the associated G/B -fibration, where B is a Borel subgroup of G , then $H^*(Y)$ is not isomorphic to $H^*(G/B \times X, \mathbf{Z})$.*

Proof. Suppose that we have proved the proposition instead for $G \times \mathbf{G}_m$ (which also is non-special) so that we have a $G \times \mathbf{G}_m$ -torsor $P \rightarrow X$ with the requisite properties. If B is a Borel subgroup of G , then $B \times \mathbf{G}_m$ is a Borel subgroup of $G \times \mathbf{G}_m$. Hence the $(G \times \mathbf{G}_m)/(B \times \mathbf{G}_m)$ -fibration associated to $P \rightarrow X$ is also the G/B -fibration associated to the G -torsor $P/\mathbf{G}_m \rightarrow X$ and we get the result for G .

Hence we may assume that G has a centrally embedded $\mathbf{G}_m$. We let X be a high degree approximation to BG as in Theorem 1.3. Then the induced morphism $Y \rightarrow G/B \times_G EG$ is also highly (cohomologically) connected. Now, $G/B \times_G EG$ is isomorphic to $EG/B = B(B) \sim BT$, where T is a maximal torus of G . This means that $H^*(G/B \times_G EG, \mathbf{Z}) = H^*(BT, \mathbf{Z})$ and $H^*(BT, \mathbf{Z})$ is the symmetric algebra on the cocharacter group of T . In particular it is torsion free so if $H^*(BG, \mathbf{Z})$ has torsion then we do not have an isomorphism $H^*(BT) \cong H^*(G/B \times BT, \mathbf{Z})$. As X can be chosen to be an arbitrarily high degree approximation if $H^*(BG, \mathbf{Z})$ has torsion we can find an X fulfilling the conditions of the proposition. However, by [SC58, Exp 5: Thm 3, Thm 4] $H^*(BG, \mathbf{Z})$ has torsion when G is non-special. $\square$

Remark: i) In many cases one can remove the condition that G be reductive. For instance, in characteristic zero any linear connected algebraic group G contains a complement to its unipotent radical U , i.e., a subgroup $H \subseteq G$ which maps isomorphically to G/U . We can then construct an H -torsor as in the proposition and extend it to a G -torsor through the inclusion $H \subseteq G$. We then have that the associated G/B -fibration is isomorphic to the H/B' -fibration associated to the original torsor.

ii) In some cases at least one can prove the proposition through the Atiyah-Hirzebruch result. For instance for SO_n one can look at the diagonally embedded $(\mathbf{Z}/2)^{n-1} \subseteq \mathrm{SO}_n$. Its classifying space detects all the torsion of $H^*(\mathrm{BSO}_n, \mathbf{Z})$ so getting a $(\mathbf{Z}/2)^{n-1} \times \mathbf{G}_m$ -torsor will give the proposition. Similarly, one can use the $\mathbf{Z}/n$ Heisenberg group and its embedding into $\mathrm{SL}_n(\mathbf{C})$ giving an embedding of $(\mathbf{Z}/n)^2 \subseteq \mathrm{PSL}_n(\mathbf{C})$ which detects the torsion class of $H^3(\mathrm{PSL}_n(\mathbf{C}), \mathbf{Z})$. In characteristic p and for a reductive group G defined over a finite field one can use $G(\mathbf{F})$ for a large enough finite field. This inclusion does not necessarily lift to a homomorphism $G(\mathbf{F}) \rightarrow G(\mathbf{C})$ (in fact typically it does not) and only gives a map from $BG(\mathbf{F})$ to the profinite completion (away from p) of $BG(\mathbf{C})$ which is not enough for our purposes (we would really need a group homomorphism $G(\mathbf{F}) \rightarrow G(\mathbf{C})$).

iii) Using the refinement discussed in the fifth part of the remark after Theorem 1.3 one can get a more precise result in that one can also demand that $H^*(Y, \mathbf{Z})$ is torsion free and as $H^*(X, \mathbf{Z})$ contains torsion it is clear that $H^*(Y, \mathbf{Z})$ is not isomorphic to $H^*(X \times G/B, \mathbf{Z})$.

Though it seems unlikely that the approximation theorem can be used to obtain any substantial properties of BG one can apply it to reprove a result of Deligne (see [De74, Thm 9.1.1]) on the rational cohomology of BG . Recall that BG can be realised as (the geometric realisation of) a simplicial scheme and hence its rational cohomology has a natural mixed Hodge structure. The result of Deligne implies that the mixed Hodge structure on $H^i(BG, \mathbf{Q})$ is pure of weight i . This however follows directly from the approximation theorem and the fact that the i 'th cohomology group of a smooth and proper variety is pure of weight i . Deligne's result is more precise in that he proves that $H^i(BG, \mathbf{Q})$ is zero for odd i and purely of type $(i/2, i/2)$ for even i . This, however, follows from the weaker result: We have the Leray spectral sequence $H^i(BG, \mathbf{Q}) \otimes H^j(G/B, \mathbf{Q}) \implies H^{i+j}(BB, \mathbf{Q})$ which is a spectral sequence of mixed Hodge structures whose E_2 -term is pure (of weight equal to the total degree). Hence, the spectral sequence degenerates for weight reasons and we get as a consequence the, *splitting principle*, that $H^*(BG, \mathbf{Q}) \rightarrow H^*(BB, \mathbf{Q})$ is injective which reduces the stronger statement to a computation of the mixed Hodge structure on $H^*(BB, \mathbf{Q}) = H^*(BT, \mathbf{Q})$. Deligne goes in the other direction, using the degeneration of the spectral sequence to get the splitting principle and hence the purity of $H^*(BG, \mathbf{Q})$. The degeneration of the spectral sequence is also equivalent to the fact that $H^*(BT, \mathbf{Q}) \rightarrow H^*(G/B, \mathbf{Q})$ is surjective which was also proved by Grothendieck (see [SC58, Exp 5: Cor 4, Lemme 10]). In any case the approximation theorem gives arguably a quite natural explanation for the fact that $H^*(BG, \mathbf{Q})$ is pure.

For our main application of the theorem we need to recall some facts. For a field $\mathbf{k}$ one defines $K_0(\mathrm{Spc}_{\mathbf{k}})$ as the group generated by elements $[X]$, X an algebraic $\mathbf{k}$ -variety, such that isomorphic varieties give the same element and $[X] - [Y] = [X \setminus Y]$ for a closed subvariety $Y \subseteq X$. It has a ring structure characterised by $[X][Y] = [X \times Y]$ and one defines a dimension filtration on the localisation $K_0(\mathrm{Spc}_{\mathbf{k}})[\mathbb{L}^{-1}]$, with $\mathbb{L} := [\mathbf{A}^1]$, where $\mathrm{Fil}^n K_0(\mathrm{Spc}_{\mathbf{k}})[\mathbb{L}^{-1}]$ is generated by elements of the form $[X]\mathbb{L}^{-i}$ for which $\dim X \leq n+i$. Then $\widehat{K}_0(\mathrm{Spc}_{\mathbf{k}})$ is the completion of $K_0(\mathrm{Spc}_{\mathbf{k}})[\mathbb{L}^{-1}]$ with respect to this filtration. If G is a special algebraic group it follows easily that for any algebraic G -torsor $P \rightarrow X$ we have $[P] = [G][X]$ in $K_0(\mathrm{Spc}_{\mathbf{k}})$. We aim to use Proposition 2.1 to give a (partial) converse to this fact.

Theorem 2.2 *For every connected non-special linear algebraic G (over the field of complex numbers) there is an algebraic G -torsor $P \rightarrow X$ such that $[P] \neq [G][X] \in \widehat{K}_0(\mathrm{Spc}_{\mathbf{k}})$.*

Proof. As G is non-special and its unipotent radical U is special (cf., [SC58, Exp 1, Prop 14]) we get that G/U is non-special as extensions of special groups are special (cf., [SC58, Exp 1, Lemme 6]) and we may assume that G is reductive. In [Ek09a, Prop. 3.2] an invariant of elements of $\widehat{K}_0(\mathrm{Spc}_{\mathbf{k}})$ is defined whose value on $[X]$ for X smooth and proper gives information equivalent to the isomorphism class of $H^*(X, \mathbf{Z})$ as a graded group. It is also proved (cf., [Ek09a, Lemma 3.8]) that if $[P] = [G][X]$ in $\widehat{K}_0(\mathrm{Spc}_{\mathbf{k}})$ for all G -torsors, then $[Y] = [G/B][X] = [G/B \times X]$ for all G/B -fibrations $Y \rightarrow X$ with structure group G . However, if we had $[Y] = [G/B \times X]$ for a smooth and proper X , then by what was just recalled (and the fact that G/B is smooth and proper) we would get $H^*(Y, \mathbf{Z}) \cong H^*(G/B \times X, \mathbf{Z})$ which by Proposition 2.1 is not always true. $\square$

Remark: i) It is not difficult to extend the theorem to non-connected groups (which are always non-special). In that case however $[P] = [G][B]$ should essentially never be true, for instance if G is finite it is true only if the torsor is trivial.

ii) Note that having a base field of characteristic zero is (currently) necessary: The invariant of [Ek09a] is defined using resolution of singularities.

iii) That invariant is additive and takes value in a torsion free abelian group so that we get not only that $[P] \neq [G][X]$ but also that the difference $[P] - [G][X]$ has infinite order.

References

- [AH62] M. F. Atiyah and F. Hirzebruch, *Analytic cycles on complex manifolds*, Topology **1** (1962), 25–45.
- [DGMS] P. Deligne, P. Griffiths, J. Morgan, and D. Sullivan, *Real homotopy theory of Kähler manifolds*, Invent. Math. **29** (1975), no. 3, 245–274.
- [De74] P. Deligne, *Theorie de Hodge, III*, Publ. Math. IHES (1974), no. 44, 5–78.
- [De77] ———, *Cohomologie étale: les points de départ*, Sem. de Géom. alg. du Bois-Marie SGA4 $\frac{1}{2}$, SLN, no. 569, Springer-Verlag, 1977, pp. 4–75.
- [Ek09a] T. Ekedahl, *The Grothendieck group of algebraic stacks*, 2009, [arXiv:math/0903.3143](#).
- [Ek86] T. Ekedahl, *Two examples of smooth projective varieties with nonzero Massey products*, Algebra, algebraic topology and their interactions (Stockholm, 1983), Lecture Notes in Math., vol. 1183, Springer, Berlin, 1986, pp. 128–132.
- [Ki84] F. C. Kirwan, *Cohomology of quotients in symplectic and algebraic geometry*, Mathematical notes, vol. 31, Princeton University Press, Princeton, NJ, 1984.
- [Mi63] J. Milnor, *Morse theory*, Based on lecture notes by M. Spivak and R. Wells. Annals of Mathematics Studies, No. 51, Princeton University Press, Princeton, N.J., 1963.
- [SC58] *Séminaire C. Chevalley; 2e année: 1958. Anneaux de Chow et applications*, Secrétariat mathématique, 11 rue Pierre Curie, Paris, 1958.
- [Se50] A. Seidenberg, *The hyperplane sections of normal varieties*, Trans. Amer. Math. Soc. **69** (1950), 357–386.
- [Se58] J.-P. Serre, *Sur la topologie des variétés algébriques en caractéristique p* , Symposium internacional de topología algebraica, Universidad Nacional Autónoma de México and UNESCO, Mexico City, 1958, pp. 24–53.

MATEMATISKA INSTITUTIONEN, STOCKHOLMS UNIVERSITET, SE-106 91 STOCKHOLM, SWEDEN

E-mail address: teke@math.su.se